



NATIONAL COMMISSIONER OF THE ICELANDIC POLICE

DEPARTMENT OF CIVIL PROTECTION AND EMERGENCY MANAGEMENT



THE SCIENTIFIC ADVISORY BOARD OF THE ICELANDIC CIVIL PROTECTION

Date: 20.01.2015 **Time:** 09:30 **Location:** Crisis Coordination Centre, Skogarhlid.

Regarding: Volcanic activity in the Bardarbunga system.

Attending: Scientists from Icelandic Met Office and the Institute of Earth Sciences University of Iceland along with representatives from the Icelandic Civil Protection, Directorate of Health and The Environment Agency of Iceland.

Main points

- Volcanic eruption in Holuhraun
- Air quality
- Scenarios

Notes

- Insubstantial changes have been in the volcanic eruption in Holuhraun over the last few weeks.
- Seismic activity in Bardarbunga continues to be strong. Over the last few weeks it has though been quite weaker than in the first months of the event. The strongest earthquake since the last meeting of the Advisory Board on last Friday was measured M4,8 on Sunday January 18th at 22:27. Another earthquake of magnitude M4,7 was detected this morning at 10:32. Sixteen earthquakes stronger than M4,0 were detected over the period and about 25 earthquakes between M3,0-4,0. In total around 230 earthquakes have been detected around the caldera since last Friday.
- Around 60 earthquakes were detected in the dyke during the same period, all of them smaller than M2,0.
- Around 15 earthquakes were detected in Tungnafellsjokull glacier since last Friday, all smaller than M2,5.
- GPS measurements near northern Vatnajokull glacier show continuing slow deflation towards Bardarbunga.
- A number of monitoring equipment on and around Vatnajokull glacier are not transmitting data. A repair mission is heading towards the glacier today but weather conditions on this time of the year are very rough.
- High values of sulphuric dioxide are still being recorded. Around 1800 µg/m³ SO₂ were recorded last Friday in Hofn in Hornarfjordur and around 1750 µg/m³ in Reykjahlid last Sunday.

Air quality:

- Today (Tuesday) the pollution from the Holuhraun eruption will move towards northwest and later north with calm winds. Tomorrow (Wednesday) calm winds are expected early on with gas pollution around the volcano and west of it. In the afternoon the winds will slowly increase and the pollution will be carried towards north.
- The Icelandic Met Office provides two-day forecasts on gas dispersion from the eruptive site in Holuhraun. Most reliable are the forecast maps approved by meteorologist on duty, see [Gas forecast](#). And although still being developed further, an automatic forecast, see [Gas model](#), is also available (trial run, see [disclaimer](#)).
- Measurements of air quality can be found on the webpage www.airquality.is Data from handheld gas monitors, spread around the country, can also be found on that page

- Instructions:



NATIONAL COMMISSIONER OF THE ICELANDIC POLICE

DEPARTMENT OF CIVIL PROTECTION AND EMERGENCY MANAGEMENT



- People who feel discomfort are advised to stay indoors, close their windows, turn up the heat and turn off air conditioning. Use periods of good air quality to ventilate the house. People experiencing adverse effects should be in immediate contact with their healthcare centre. Measurements of air quality can be found on the webpage www.airquality.is The Meteorological Office issues forecast on its web-page and warnings if conditions change to the worse.
 - Instructions from [The Environment Agency of Iceland](#) and [Chief Epidemiologist](#) can be found on their web-sites.
 - Check the Icelandic Met Office forecasts for sulphuric gas dispersion on the web as described above.
 - Handheld meters have been distributed around the country for SO₂ measurements three times a day.
 - Information and any questions on air pollution can be sent to The Environment Agency through the email gos@ust.is. The Environment Agency is especially looking for information from people who have been in contact with high concentrations of gas; where they were, at what time it happened, how the gas cloud looked (colour and thickness of the cloud) and how they were affected by it.
- The volcanic eruption has now been going on for over four months, the lava flow is still great in Holuhraun and the rate of the subsidence of the Bardarbunga caldera is still significant. Three scenarios are considered most likely:
 - The eruption in Holuhraun continues until the subsidence of the Bardarbunga caldera stops. The eruption can still go on for many months.
 - The volcanic fissure may lengthen southwards under Dyngjajokull, resulting in a jokulhlaup and an ash-producing eruption. It is also possible that eruptive fissures could develop in another location under the glacier. If such an eruption would be prolonged it could eventually produce a lava flow.
 - Volcanic eruption in the Bardarbunga caldera. Such an eruption would melt large quantities of ice, leading to a major jokulhlaup, accompanied by ash fall.
- Other scenarios cannot be excluded.
- **From the Icelandic Met Office:** The Aviation Colour Code for Bardarbunga remains at 'orange'.
 - The next meeting will be held on Friday 23rd of January 2015.

The National Commissioner of the Icelandic Police, Department of Civil Protection and Emergency Management
www.avd.is/en Twitter: [@almannavarnir](https://twitter.com/almannavarnir)